

AD-A123 916

WAVE PROPAGATION-NONLINEAR BOUNDARY VALUE PROBLEMS(U)
MIAMI UNIV CORAL GABLES FLA DEPT OF MATHEMATICS
R A GOLDSTEIN 25 JAN 82 N00014-78-C-0608

1/1

UNCLASSIFIED

F/G 20/9

NL




END

FILED

F

DEC



ADA 123916



University of ~~Florida~~
Coral Gables, Florida 33124

DEPARTMENT OF MATHEMATICS

P.O. Box 249085

Coleman Hall, University Center

Annual Report

For

Navy Contract N00014-78-C-0608

(UM A/C BR6537)

"Wave Propagation-Nonlinear Boundary Value Problems"

Principal Investigator: R.A. Goldstein

25 January 1982

APPROVED FOR PUBLIC RELEASE
DISTRIBUTION UNLIMITED

DTIC
ELECTE
JAN 28 1983
S D D

DTIC FILE COPY

83 01 23 032

Work was completed on a paper entitled "Perturbed Bifurcation of Stationary Striations in a Contaminated, Non-Uniform Plasma." It was accepted by SIAM APPLIED MATH and is currently in press. The paper is attached.

On going research in collaboration with J. Magnan (Northwestern) continues with respect to moving situations or modulated travelling waves in reaction diffusion systems.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

Copy
/New Access
2

Moung Striations

Problem

$$\frac{\partial N}{\partial t} = D_- \frac{\partial^2 N}{\partial z^2} + \frac{\mu_+}{\mu_-} \frac{J}{e} \left(\frac{P}{N^2} \frac{\partial N}{\partial z} - \frac{1}{N} \frac{\partial P}{\partial z} \right) + F,$$

$$\frac{\partial P}{\partial t} = D_+ \frac{\partial^2 P}{\partial z^2} + \frac{\mu_+}{\mu_-} \frac{J}{e} \left(\frac{P}{N^2} \frac{\partial N}{\partial z} - \frac{1}{N} \frac{\partial P}{\partial z} \right) + G,$$

$$\frac{\partial M}{\partial t} = D^* \frac{\partial^2 M}{\partial z^2} + H,$$

where it has been assumed that $\mu_- N \gg \mu_+ P$

B.C. (see Fig 1)

$$N(0) - \alpha \frac{\partial N(0)}{\partial z} = A, \quad N(L) + \alpha \frac{\partial N(L)}{\partial z} = A$$

$$P(0) - \beta \frac{\partial P(0)}{\partial z} = B, \quad P(L) + \beta \frac{\partial P(L)}{\partial z} = B$$

$$M(0) - \gamma \frac{\partial M(0)}{\partial z} = C, \quad M(L) + \gamma \frac{\partial M(L)}{\partial z} = C$$


I.C.

$$N(z, 0) = N_0$$

$$P(z, 0) = P_0$$

$$M(z, 0) = M_0$$

Fig 1



Scaled equations

$$\frac{\partial n}{\partial \tau} = \epsilon \frac{\partial^2 n}{\partial \xi^2} + \frac{p}{n^2} \frac{\partial n}{\partial \xi} - \frac{1}{n} \frac{\partial p}{\partial \xi} + f$$

$$\frac{\partial p}{\partial \tau} = \epsilon \delta \frac{\partial^2 p}{\partial \xi^2} + \frac{p}{n^2} \frac{\partial n}{\partial \xi} - \frac{1}{n} \frac{\partial p}{\partial \xi} + g$$

$$\frac{\partial m}{\partial \tau} = \epsilon \theta \frac{\partial^2 m}{\partial \xi^2} + h$$

B.C.

$$n(0, \tau) - \epsilon \frac{\partial n(0, \tau)}{\partial \xi} = a = n(l, \tau) + \epsilon \frac{\partial n(l, \tau)}{\partial \xi}$$

$$p(0, \tau) + \epsilon \frac{\partial p(0, \tau)}{\partial \xi} = b = p(l, \tau) + \epsilon \frac{\partial p(l, \tau)}{\partial \xi}$$

$$m(0, \tau) - \epsilon \frac{\partial m(0, \tau)}{\partial \xi} = c = m(l, \tau) + \epsilon \frac{\partial m(l, \tau)}{\partial \xi}$$

I.C.

$$n(\frac{1}{2}, 0) = 1$$

$$p(\frac{1}{2}, 0) = 1$$

$$m(\frac{1}{2}, 0) = 1$$

where

$$z = \left(\frac{L^2}{D_-} \right) \frac{\mu_+}{\mu_-} \frac{J}{e N_0} \frac{1}{\xi} ,$$

$$t = \frac{L^2}{D_-} \tau ,$$

$$\epsilon = \left(\frac{\frac{D_- N_0}{L}}{\frac{\mu_+}{\mu_-} \frac{J}{e}} \right)^2 = \left(\frac{\Gamma_0}{\Gamma_+} \right)^2 ,$$

$$N = n N_0 , P = p N_0 , M = m M_0 ,$$

$$A = a N_0 , B = b N_0 , C = c M_0 ,$$

$$\alpha = \beta = \gamma = \sqrt{\epsilon} L ,$$

$$\delta = \left(\frac{D_+}{D_-} \right)^2 , \theta = \left(\frac{D^*}{D_-} \right)^2 , \ell = \sqrt{\epsilon} .$$

Xform to homogeneous B.C.

$$\text{let } n = n' + a$$

$$p = p' + b$$

$$m = m' + c$$

substitute into the scaled equations and drop the primes.

$$\frac{\partial n}{\partial \tau} = \epsilon \frac{\partial^2 n}{\partial \xi^2} + \frac{(p+b)}{(n+a)^2} \frac{\partial n}{\partial \xi} - \frac{1}{(n+a)} \frac{\partial p}{\partial \xi} + f$$

$$\frac{\partial p}{\partial \tau} = \epsilon \delta \frac{\partial^2 p}{\partial \xi^2} + \frac{(p+b)}{(n+a)^2} \frac{\partial n}{\partial \xi} - \frac{1}{(n+a)} \frac{\partial p}{\partial \xi} + g$$

$$\frac{\partial m}{\partial \tau} = \epsilon \theta \frac{\partial^2 m}{\partial \xi^2} + h$$

BC:

$$n(0, \tau) - \epsilon \frac{\partial n(0, \tau)}{\partial \xi} = 0 = n(l, \tau) + \epsilon \frac{\partial n(l, \tau)}{\partial \xi}$$

$$p(0, \tau) - \epsilon \frac{\partial p(0, \tau)}{\partial \xi} = 0 = p(l, \tau) + \epsilon \frac{\partial p(l, \tau)}{\partial \xi}$$

$$m(0, \tau) - \epsilon \frac{\partial m(0, \tau)}{\partial \xi} = 0 = m(l, \tau) + \epsilon \frac{\partial m(l, \tau)}{\partial \xi}$$

IC:

$$n(\xi, 0) = 1 - a$$

$$p(\xi, 0) = 1 - b$$

$$m(\xi, 0) = 1 - c$$

Try a solution

$$\text{let } u(\xi, \tau) = e^{-\frac{\phi(\xi, \tau)}{2\epsilon}} x(\xi, \tau, \epsilon),$$

$$p(\xi, \tau) = e^{-\frac{\phi(\xi, \tau)}{2\epsilon}} y(\xi, \tau, \epsilon),$$

$$w(\xi, \tau) = e^{-\frac{\phi(\xi, \tau)}{2\epsilon}} z(\xi, \tau, \epsilon),$$

substitute into equations and get
to order $\frac{1}{\epsilon}$:

$$\phi_\tau + \frac{1}{2} \phi_{\xi\xi}^2 = 0$$

with solution

$$\phi(\xi, \tau) = \frac{1}{2} \left(\frac{\xi - \xi_0}{\tau} \right)^2$$

$\begin{pmatrix} x_0 \\ y_0 \\ z_0 \end{pmatrix}$ will satisfy the 1st order unperturbed problem (hence a travelling wave)

We will seek "modulated" travelling wave solutions